

GOËMAR®

BM 86

5-0-0

FOLIAR NUTRITIONAL FORMULATION

GUARANTEED ANALYSIS:

Total Nitrogen (N).....	5.0%
5.00% Water Soluble Nitrogen	
Magnesium (Mg).....	2.4%
2.4% Water soluble magnesium (Mg)	
Sulfur (S) combined.....	3.2%
Boron (B).....	2.0%
Molybdenum (Mo).....	0.02%
Sodium (Na).....	0.6%

Derived from: Urea, Magnesium sulfate, boric acid, sodium molybdate.

DISTRIBUTED BY:

Agrimar, Corporation
5634 Atlanta Highway
Suite 200
Flowery Branch, GA 30542
800-638-6673

Net weight: 26.45 lbs. (12.0 kg) Net content: 2.5 gal. (9.46 liters)
Weight per gallon: 10.6 lbs. (4.8 kgs)

CAUTION: KEEP OUT OF REACH OF CHILDREN

CAUTION: The application of fertilizing materials containing molybdenum (Mo) may result in crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

CAUTION: This product contains boron, therefore must only be used on crops listed on this label at the recommended rates and according to the directions for use. The use of this fertilizer material on any other crops may result in serious crop injury.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

**Product of France
FRONT JUG PANEL**

VEGETABLE CROPS

Fruit-bearing vegetables: For fruit-bearing vegetable crops not specifically listed on label, apply Goëmar BM 86 at pre-bloom, bloom and every one to two weeks thereafter at a rate of 3 pints to 2 quarts per acre.

TOMATOES, PEPPERS, CUCURBITS AND OTHER FRUITING VEGETABLES: Dosage per acre per application: 3 pints. Three applications: 1) at first flower. 2) 7 - 14 days later. 3) 7 - 14 days later.

POTATOES (Irish): Dosage per acre per application: 3 pints. Two applications. 1) 100% sprouting. 2) 15 days after first treatment.

SNAP BEANS, PEAS AND OTHER LEGUMES: Dosage per acre per application: 2 quarts. Two applications. 1) At 1st bloom. 2) At first pod.

ROW & NUT CROPS

COTTON: Dosage per acre per application: 1 quart. One application: 1) At early to mid bloom.

PEANUTS: Dosage per acre per application: 1 quart. Two applications: 1) At pegging. 2) 10 - 14 days after.

ALMONDS & PISTACHIOS: Dosage per acre per application: 2 quarts. Two applications: 1) Pre-bloom to early bloom. 2) Full bloom to petal fall. OR One 3 qt/acre application @ full bloom.

ENGLISH WALNUTS AND PECANS
Dosage per acre per application: Two quarts. Two applications: 1) At early pistillate bloom. 2) At full pistillate bloom.

SUGAR BEET: 3 pints per acre applied at 8-10 leaf stage followed by 3 pints per acre 14-21 days later.

SUNFLOWER: 2.5 pints per acre applied at 3 pairs of true leaves to flower bud appearance.

FRUIT CROPS

Apply Goëmar BM 86 to most fruit crops at pre-bloom, full bloom and soon after fruit set at a rate of 3 pints to 2 quarts per acre.

APPLES & PEARS: Dosage per acre per application: 3 pints. Three applications: 1) Pre-bloom. 2) Early king bloom (for apples)/bloom (for pears). 3) Post bloom (petal fall - fruit set).

GRAPES (Raisin, Table and Wine): Three applications: 1) 12 - 18 inch shoot growth, 1 quart per acre. 2) At bloom stage, 1 quart per acre. 3) After fruit set, 1 quart per acre.

Concord, Niagara and other American cultivars: Two applications 1) Early bloom, 2 quarts per acre. 2) Full bloom, 2 quarts per acre.

STONE FRUIT (Peaches, Nectarines, Apricots, Plums): Dosage per acre per application: 3 pints. Three applications: 1) Early bloom. 2) Petal fall. 3) 10 - 20 days later (dependent upon variety).

CHERRIES: Dosage per acre per application: 2 quarts. Three applications: 1) At first white flower. 2) At petal fall. 3) At shuck fall.

STRAWBERRIES: Dosage per acre per application: 1 quart. 1) Within 14 days of transplant. 2) At first bloom. 3) 14 - 21 days later. Subsequent applications: Every 14 - 21 days thereafter.

BLUEBERRY, CRANBERRY AND CANE BERRY: Dosage per acre per application: 1 quart. Three applications: 1) At first bloom. 2) At full bloom. 3) At fruit set.

CAUTION: The application of fertilizing materials containing molybdenum (Mo) may result in crops containing levels of molybdenum (Mo), which are toxic to ruminant animals.

CAUTION: Goëmar BM 86 contains boron, therefore must only be used on crops listed on this label at the recommended rates and according to the directions for use. The use of this fertilizer material on any other crops may result in serious crop injury.

For information on specific crop recommendations, please call 800-638-6673.